

Proposal 5

A Longitudinal-Transverse Separation of Amplitudes for the $S_{11}(1535)$ Resonance

The N^* -Collaboration

G.Adams, R.Arndt, V.Burkert, C.Carlson, D.Day, L. Dennis, T.Donoghue, D.Doughty, S. Dytman, H.Funsten, M.Gai, L.Ghedira, K.Giovanetti, D. Isenhower, D. Jenkins, D.Joyce, D.Keane, K. Kemper, W.Kim, J. Lieb, M.Manley, J.McCarthy, B.Mecking, M.Mestayer, R.Minehart, N.Mukhopadhyay, B.Niczyporuk, O.Rondon-Aramayo, D.Roper, M. Sadler, R.Sealock, E.Smith, P.Stoler, C.Stronach, S.Thornton, H.Weber, A.Yegneswaran

Spokesmen: K. Giovanetti and S. Dytman

October 30, 1989

ABSTRACT

We propose to measure the longitudinal contribution to the electroproduction of the $S_{11}(1535)$ resonance as a function of Q^2 . The properties of the S_{11} form factor provide excellent evidence for the breakdown of $SU(6)$ symmetry. We propose this case because in addition to its interesting physics properties, the S_{11} has a large production cross section and is easily identified through the ηp decay channel. Because all resonances are predominantly excited by transverse photons, an absolute accuracy of a few percent will be required. Given the design goals of the CEBAF electron beam and the proposed CLAS detector we expect to be able to measure total cross sections with a few percent accuracy and extract the ratio R (σ_L/σ_T) with an error of ± 0.05 . With such stringent requirements, this experiment should run in the 2nd round of CLAS experiments.